



Complete Summary

TITLE

Chronic stable coronary artery disease: percentage of patients who were screened for diabetes.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement. Clinical performance measures. Chronic stable coronary artery disease. Chicago (IL): American Medical Association (AMA); 2003. 8 p.

Brief Abstract

DESCRIPTION

This measure assesses the percentage of patients with chronic stable coronary artery disease (CAD) who were screened for diabetes.

RATIONALE

Screening for diabetes is recommended in patients who are considered high risk (e.g., coronary artery disease [CAD]) (ADA, 2003; ACE, 2002; AACE, 2003).

PRIMARY CLINICAL COMPONENT

Coronary artery disease; screening for diabetes

DENOMINATOR DESCRIPTION

All patients with coronary artery disease (CAD) who do not have documented diabetes

NUMERATOR DESCRIPTION

The number of patients from the denominator who were screened for diabetes

Evidence Supporting the Measure

PRIMARY MEASURE DOMAIN

Process

SECONDARY MEASURE DOMAIN

Not applicable

EVIDENCE SUPPORTING THE MEASURE

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence
One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

NATIONAL GUIDELINE CLEARINGHOUSE LINK

- [ACC/AHA 2002 guideline update for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines \(Committee to Update the 1999 Guidelines for the Management of Patients With Chronic Stable Angina\).](#)

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Wide variation in quality for the performance measured

EVIDENCE SUPPORTING NEED FOR THE MEASURE

American College of Endocrinology Consensus Statement on guidelines for glycemic control. Endocr Pract 2002;8(Suppl 1):6-11.

American Diabetes Association. Clinical practice recommendations 2003. Screening for type 2 diabetes (position statement). Diabetes Care 2003;26(suppl 1):21-4.

Communication from American Association of Clinical Endocrinologists. January 2003.

State of Use of the Measure

STATE OF USE

Pilot testing

CURRENT USE

Internal quality improvement

Application of Measure in its Current Use

CARE SETTING

Ambulatory Care

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Individual Clinicians

TARGET POPULATION AGE

Patients of all ages with the diagnosis of chronic stable coronary artery disease

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Not applicable

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

- Approximately 13 million Americans are living with coronary artery disease (CAD).
- More than 1 million Americans had a new or recurrent coronary attack in 2001.

EVIDENCE FOR INCIDENCE/PREVALENCE

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

ASSOCIATION WITH VULNERABLE POPULATIONS

Not applicable

BURDEN OF ILLNESS

- Chronic stable coronary artery disease (CAD) is the leading cause of mortality in the United States, accounting for almost 1 in 5 deaths.
- For individuals with CAD, the risk of another heart attack, stroke, and other serious complications is substantial.

EVIDENCE FOR BURDEN OF ILLNESS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

UTILIZATION

Within the past 2 decades, the number of short-stay hospital discharges for individuals with coronary artery disease (CAD) increased by almost 18%.

EVIDENCE FOR UTILIZATION

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

COSTS

The total annual cost of coronary artery disease (CAD) in the United States is approximately \$130 billion.

EVIDENCE FOR COSTS

American Heart Association. Heart disease and stroke statistics - 2003 update. Dallas (TX): American Heart Association; 2002. 46 p.

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Living with Illness

IOM DOMAIN

Effectiveness

Data Collection for the Measure

CASE FINDING

Users of care only

DESCRIPTION OF CASE FINDING

This performance measure is designed for prospective data collection in the office-based practice only. The measurement period may begin with the date of the most recent office visit, regardless of the diagnosis at that visit, and the data collection continues until 12 months are completed.

DENOMINATOR SAMPLING FRAME

Patients associated with provider

DENOMINATOR (INDEX) EVENT

Clinical Condition

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients with coronary artery disease (CAD) who do not have documented diabetes

Exclusions

Patients with documented diabetes

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

All patients who were screened for diabetes*

*Screening for diabetes is usually done by fasting blood glucose or 2-hour glucose tolerance testing. Clinical recommendations indicate screening should be considered at 3-year intervals.

Exclusions

None

DENOMINATOR TIME WINDOW

Time window follows index event

NUMERATOR TIME WINDOW

Fixed time period

DATA SOURCE

Medical record

LEVEL OF DETERMINATION OF QUALITY

Individual Case

PRE-EXISTING INSTRUMENT USED

None

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a higher score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

Unspecified

Identifying Information

ORIGINAL TITLE

Screening for diabetes.

MEASURE COLLECTION

[The Physician Consortium for Performance Improvement Measurement Sets](#)

MEASURE SET NAME

[American College of Cardiology, American Heart Association, and Physician Consortium for Performance Improvement: Chronic Stable Coronary Artery Disease Core Physician Performance Measurement Set](#)

SUBMITTER

American Medical Association on behalf of the American College of Cardiology, the American Heart Association, and the Physician Consortium for Performance Improvement

DEVELOPER

American College of Cardiology
American Heart Association
Physician Consortium for Performance Improvement

ADAPTATION

Measure was not adapted from another source.

RELEASE DATE

2003 Jan

MEASURE STATUS

This is the current release of the measure.

SOURCE(S)

American College of Cardiology, American Heart Association, Physician Consortium for Performance Improvement. Clinical performance measures. Chronic stable coronary artery disease. Chicago (IL): American Medical Association (AMA); 2003. 8 p.

MEASURE AVAILABILITY

The individual measure, "Screening for Diabetes," is published in the "Chronic Stable Coronary Artery Disease Core Physician Performance Measurement Set." This document is available from the American Medical Association (AMA) Division of Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.

For further information, please contact AMA staff by e-mail at cqi@ama-assn.org.

COMPANION DOCUMENTS

The following are available:

- Physician Consortium for Performance Improvement. Introduction to physician performance measurement sets. Tools developed by physicians for physicians. Chicago (IL): American Medical Association (AMA); 2001 Oct. 21 p. This document is available from the American Medical Association (AMA) Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.
- Physician Consortium for Performance Improvement. Principles for performance measurement in health care. A consensus statement. [online]. Chicago (IL): American Medical Association (AMA), Joint Commission on the Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); [3 p]. This document is available from the AMA Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.
- Physician Consortium for Performance Improvement. Desirable attributes of performance measures. A consensus statement. [online]. American Medical Association (AMA), Joint Commission on Accreditation of Healthcare Organizations (JCAHO), National Committee for Quality Assurance (NCQA); 1999 Apr 19 [cited 2002 Jun 19]. [5 p]. This document is available from the AMA Clinical Quality Improvement Unit Web site: www.ama-assn.org/go/quality.

For further information, please contact AMA staff by e-mail at cqi@ama-assn.org.

NQMC STATUS

This NQMC summary was completed by ECRI on September 26, 2003.

COPYRIGHT STATEMENT

This Physician Performance Measurement Set (PPMS) was developed by the Physician Consortium for Performance Improvement (The Consortium) to facilitate quality improvement activities by physicians. The performance measures contained in this PPMS are not clinical guidelines and do not establish a standard of medical care. This PPMS is intended to assist physicians in enhancing quality of care and is not intended for comparing individual physicians to each other or for individual physician accountability by comparing physician performance against the measure or guideline. The Consortium has not tested this PPMS.

This PPMS is subject to review and may be revised or rescinded at any time by The Consortium. The PPMS may not be altered without the prior written approval of The Consortium. A PPMS developed by The Consortium, while copyrighted, can be reproduced and distributed without modification for noncommercial purposes. Any other use is subject to the approval of The Consortium. Neither The Consortium nor its members shall be responsible for any use of this PPMS.

©2003 American Medical Association

© 2004 National Quality Measures Clearinghouse

Date Modified: 11/1/2004

The logo for FIRSTGOV, featuring the word "FIRST" in blue and "GOV" in red, with a stylized red and blue graphic above the "I" in "FIRST".

