



Complete Summary

TITLE

Postoperative wound dehiscence (area-level): rate of reclosure of abdominal wall per 100,000 population.

SOURCE(S)

AHRQ quality indicators. Guide to patient safety indicators [revision 1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2003 May 28. 143 p. (AHRQ Pub; no. 03-R203).

Brief Abstract

DESCRIPTION

This measure assesses the number of cases of reclosure of postoperative disruption of abdominal wall per 100,000 population.

RATIONALE

Patient safety is an issue of major national interest. Policymakers, providers, and consumers have made the safety of care in United States hospitals a top priority. The need to assess, monitor, track, and improve the safety of inpatient care became apparent with publication of the Institute of Medicine's series of reports describing the problem of medical errors. As our health care system becomes more complex, the possibility of significant unintended adverse effects increases.

Patient Safety Indicators (PSIs), which are based on computerized hospital discharge abstracts from the AHRQ's Healthcare Cost and Utilization Project (HCUP), can be used to better prioritize and evaluate local and national initiatives. Analyses of these and similar inexpensive, readily available administrative data sets may provide a screen for potential medical errors and a method for monitoring trends over time.

This indicator is intended to flag cases of wound dehiscence in patients who have undergone abdominal and pelvic surgery. This indicator is defined both on a hospital level (by including cases based on secondary diagnosis associated with the same hospitalization) (see [Postoperative wound dehiscence \(hospital-level\): rate of reclosure per 1,000 cases of abdominopelvic surgery](#)) and on an area level (by including all cases of wound dehiscence).

PRIMARY CLINICAL COMPONENT

Postoperative wound dehiscence

DENOMINATOR DESCRIPTION

Population of county or Metropolitan Statistical Area (MSA) associated with Federal Information Processing Standards (FIPS) code of patient's residence or hospital location.

NUMERATOR DESCRIPTION

Discharges with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code for reclosure of postoperative disruption of abdominal wall (5461) in any procedure field (principal or secondary) of abdominopelvic surgical discharges.

Exclude obstetrical patients in Major Diagnostic Category 14 (MDC 14).

Evidence Supporting the Measure

PRIMARY MEASURE DOMAIN

Outcome

SECONDARY MEASURE DOMAIN

Not applicable

EVIDENCE SUPPORTING THE MEASURE

A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences
One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Evidence Supporting Need for the Measure

NEED FOR THE MEASURE

Unspecified

State of Use of the Measure

STATE OF USE

Current routine use

CURRENT USE

Federal health policymaking
National health care quality reporting
State health policymaking

Application of Measure in its Current Use

CARE SETTING

Hospitals

PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Physicians
Public Health Professionals

LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED

Counties or Cities

TARGET POPULATION AGE

Unspecified

TARGET POPULATION GENDER

Either male or female

STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

Characteristics of the Primary Clinical Component

INCIDENCE/PREVALENCE

Based on the 1997 Healthcare Cost and Utilization Project (HCUP) State Inpatient Database for 19 States, the Postoperative Wound Dehiscence (Area-Level) rate was 1.36 per 100,000 population.

EVIDENCE FOR INCIDENCE/PREVALENCE

AHRQ quality indicators. Guide to patient safety indicators [revision 1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2003 May 28. 143 p. (AHRQ Pub; no. 03-R203).

ASSOCIATION WITH VULNERABLE POPULATIONS

Unspecified

BURDEN OF ILLNESS

Based on the matching analysis of the 2000 Healthcare Cost and Utilization Project (HCUP) Nationwide Inpatient Sample (NIS) data reported by Zhan and

Miller, postoperative wound dehiscence was associated with an excess length of stay of 9.42 days, excess charges of \$40,323, and an excess mortality rate of 9.63%.

EVIDENCE FOR BURDEN OF ILLNESS

Zhan C, Miller MR. Excess length of stay, charges, and mortality attributable to medical injuries during hospitalization. JAMA 2003 Oct 8;290(14):1868-74.
[PubMed](#)

UTILIZATION

See "Burden of Illness" field.

COSTS

See "Burden of Illness" field.

Institute of Medicine National Healthcare Quality Report Categories

IOM CARE NEED

Getting Better

IOM DOMAIN

Safety

Data Collection for the Measure

CASE FINDING

Both users and nonusers of care

DESCRIPTION OF CASE FINDING

Population of county or Metropolitan Statistical Area (MSA) associated with Federal Information Processing Standards (FIPS) code of patient's residence or hospital location.

DENOMINATOR SAMPLING FRAME

Geographically defined

DENOMINATOR (INDEX) EVENT

Patient Characteristic

DENOMINATOR INCLUSIONS/EXCLUSIONS

Inclusions

Population of county or Metropolitan Statistical Area (MSA) associated with Federal Information Processing Standards (FIPS) code of patient's residence or hospital location.

Exclusions

Unspecified

NUMERATOR INCLUSIONS/EXCLUSIONS

Inclusions

Discharges with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code for reclosure of postoperative disruption of abdominal wall (5461) in any procedure field (principal or secondary) of abdominopelvic surgical discharges. Refer to Appendix A of the original measure documentation for ICD-9-CM codes.

Exclusions

Exclude obstetrical patients in Major Diagnostic Category 14 (MDC 14) (Pregnancy, Childbirth and the Puerperium).

DENOMINATOR TIME WINDOW

Time window is a single point in time

NUMERATOR TIME WINDOW

Institutionalization

DATA SOURCE

Administrative data
National public health data

LEVEL OF DETERMINATION OF QUALITY

Not Individual Case

OUTCOME TYPE

Adverse Outcome

PRE-EXISTING INSTRUMENT USED

Unspecified

Computation of the Measure

SCORING

Rate

INTERPRETATION OF SCORE

Better quality is associated with a lower score

ALLOWANCE FOR PATIENT FACTORS

Unspecified

STANDARD OF COMPARISON

External comparison at a point in time
External comparison of time trends
Internal time comparison

Evaluation of Measure Properties

EXTENT OF MEASURE TESTING

The Patient Safety Indicators (PSIs) were evaluated by the project team using empirical analyses to explore the frequency and variation of the indicators, the potential bias, based on limited risk adjustment, and the relationship between indicators. The data sources used in the empirical analyses were the 1997 Florida State Inpatient Database (SID) for initial testing and development and the 1997 Healthcare Cost and Utilization Project (HCUP) State Inpatient Database for 19 States for the final empirical analyses.

All potential indicators were examined empirically by developing and conducting statistical tests for precision, bias, and relatedness of indicators. Three different estimates of hospital performance were calculated for each indicator:

1. The raw indicator rate was calculated using the number of adverse events in the numerator divided by the number of discharges in the population at risk by hospital.
2. The raw indicator was adjusted to account for differences among hospitals in age, gender, modified Diagnosis-Related Group (DRG), and comorbidities.
3. Multivariate signal extraction methods were applied to adjust for reliability by estimating the amount of "noise" (i.e., variation due to random error) relative to the amount of signal (i.e., systematic variation in hospital performance or reliability) for each indicator.

Refer to the original measure documentation for additional details.

EVIDENCE FOR RELIABILITY/VALIDITY TESTING

AHRO quality indicators. Guide to patient safety indicators [revision 1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRO); 2003 May 28. 143 p. (AHRO Pub; no. 03-R203).

Identifying Information

ORIGINAL TITLE

Postoperative wound dehiscence (area level definition).

MEASURE COLLECTION

[Agency for Healthcare Research and Quality \(AHRQ\) Quality Indicators](#)

MEASURE SET NAME

[Agency for Healthcare Research and Quality \(AHRQ\) Patient Safety Indicators](#)

DEVELOPER

Agency for Healthcare Research and Quality

INCLUDED IN

National Healthcare Quality Report (NHQR)

ADAPTATION

An indicator on this topic (9983) was originally proposed by Hannan and colleagues (1989) to target "cases that would have a higher percentage of quality of care problems than cases without the criterion, as judged by medical record review." The same code was included within a broader indicator ("adverse events and iatrogenic complications") in the Agency for Healthcare Research and Quality's (AHRQ's) original Healthcare Cost and Utilization Project (HCUP) Quality Indicators (Elixhauser et al., 1998). Iezzoni and colleagues (1994) identified an associated procedure code for reclosure of an abdominal wall dehiscence (5461), and included both codes in the Complications Screening Program. Miller and colleagues (2001) suggested the use of both codes (as "wound disruption") in the original "AHRQ Patient Safety Indicator (PSI) Algorithms and Groupings."

RELEASE DATE

2003 Mar

REVISION DATE

2003 May

MEASURE STATUS

This is the current release of this measure.

SOURCE(S)

AHRQ quality indicators. Guide to patient safety indicators [revision 1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2003 May 28. 143 p. (AHRQ Pub; no. 03-R203).

MEASURE AVAILABILITY

The individual measure, "Postoperative Wound Dehiscence (Area Level Definition)," is published in "AHRQ Quality Indicators. Guide to Patient Safety Indicators." This document is available in [Portable Document Format \(PDF\)](#) and a [zipped WordPerfect\(R\) file](#) from the [Quality Indicators](#) page at the Agency for Healthcare Research and Quality (AHRQ) Web site.

For more information, please contact the QI Support Team at support@qualityindicators.ahrq.gov.

COMPANION DOCUMENTS

The following are available:

- "AHRQ Patient Safety Quality Indicators Software (Version 2.1)" (Rockville, [MD]: AHRQ, 2003 Mar 13) and its accompanying documentation can be downloaded from the [Agency for Healthcare Research and Quality \(AHRQ\) Web site](#). (The software is available in both SAS- and SPSS-compatible formats.)
- Guidance for using the AHRQ quality indicators for hospital-level public reporting or payment. Rockville (MD): Agency for Healthcare Research and Quality; 2004 Aug. 24 p. This document is available from the [AHRQ Web site](#).
- "HCUPnet, Healthcare Cost and Utilization Project" [internet]. (Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2004 [Various pagings]). HCUPnet is available from the [AHRQ Web site](#).
- "Refinement of the HCUP Quality Indicators" (Rockville [MD]: AHRQ, 2001 May. Various pagings. [Technical review; no. 4]; AHRQ Publication No. 01-0035). This document was prepared by the UCSF-Stanford Evidence-based Practice Center for AHRQ and can be downloaded from the [AHRQ Web site](#).

NQMC STATUS

This NQMC summary was completed by ECRI on October 1, 2003. The information was verified by the measure developer on October 29, 2003.

COPYRIGHT STATEMENT

No copyright restrictions apply.

© 2004 National Quality Measures Clearinghouse

Date Modified: 11/1/2004

FIRST GOV

